



Bruna MOREIRA<sup>1,2,3</sup>, Ermelinda SILVA<sup>1,4</sup>, Alexandre GONÇALVES<sup>1,4</sup>, Frane STRIKIĆ<sup>5</sup>, Tânia C.S. PIRES<sup>1,2</sup>, Ricardo C. CALHELHA<sup>1,2</sup>, Miguel A. PRIETO<sup>3</sup>, Márcio CAROCHO<sup>1,2</sup>, Cristina CALEJA<sup>1,2</sup>, Lillian BARROS<sup>1,2\*</sup>

<sup>1</sup> Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

<sup>2</sup> Laboratório Associado para a Sustentabilidade e Tecnologia em Regiões de Montanha (SusTEC), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

<sup>3</sup> Nutrition and Bromatology Group, Department of Analytical and Food Chemistry, Faculty of Food Science and Technology, University of Vigo-Ourense Campus, E-32004 Ourense, Spain

<sup>4</sup> MORE - Collaborative Laboratory Mountains of Research, Brigantia Ecopark, 5300-358 Bragança, Portugal

<sup>5</sup> University of Split, Department of Mediterranean agriculture, Ruđera Boškovića 31, 21000, Split, Croatia

\* [lillian@ipb.pt](mailto:lillian@ipb.pt)



## Introduction

In recent years, crop diversification in agroforestry systems has been recognized as a sustainable strategy to improve land use efficiency, restore environmental balance, mitigate climate change, provide economic benefits and ensure food security. Research has shown that practices such as intercropping can significantly enhance soil quality, which in turn increases the quality of almonds. This study aimed to evaluate the nutritional and bioactive aspects of almonds to investigate the benefits resulting from crop interactions, using different almond cultivars.



## Results

The Texas cultivar had the highest crude fat content, while the AI cultivar had the highest protein content (Table 1).

Table 1. Nutritional profile of almonds produced in Croatia.

	Moisture (g/100 g fw)	Crude Fat (g/100 g fw)	Proteins (g/100 g fw)	Ash (g/100 g fw)	Carbohydrates (g/100 g fw)	Energy (Kcal/100g)
ACFD	6±0,2	47±2	12,3±0,02	3,54±0,3	30±0,1	600,5±1
ACFG	4,2±0,1	49±2,3	15±2	3,41±0,1	28±2	621,5±2,3
ACT	4,0±0,2	54±0,2	15±0,1	3,24±0,14	24±1	638±0,2
ACAI	5±0,1	48±0,4	19,2±0,4	3,73±0,1	25±1	605,4±2

ACT – Intercropped almond corresponding to the Texas cultivar; ACAI – Intercropped almond corresponding to the AI cultivar; ACFG - Intercropped almond corresponding to the Ferragnès cultivar; ACFD - Intercropped almond corresponding to the Ferraduel cultivar.

The Ferraduel cultivar showed the highest antioxidant activity for TBARS (Table 2).



## Conclusions

Overall, the study shows that interaction with other crops can improve the nutritional profile and health benefits of almonds, underlining the importance of sustainable agricultural practices.




Table 2. Results of lipid peroxidation inhibition assay (TBARS), 2,2- diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity, cytotoxic and anti-inflammatory activity of the almonds.

Antioxidant Activity (IC <sub>50</sub> values, µg/mL)					
	ACFG	ACT	ACAI	ACFD	
TBARS (IC <sub>50</sub> values, mg/mL)	0,2±0,01	0,4±0,003	0,2±0,001	0,1±0,003	
DPPH (mg/mL)	1±0,01	1,7±0,02	0,4±0,01	1,2±0,04	
Cytotoxicity (GI <sub>50</sub> values; µg/mL)					
	ACFG	ACT	ACAI	ACFD	
Tumoral cell lines	Caco2	66±3	>400	29±4	29±4
	MCF_7	93±4	>400	58±3	65,14±3
	AGS	102±3	>400	61,2±2	67,1±2
	HeLa	110,3±3	>400	55±2	62±2
Non-tumor cell line	VERO	161±2	>400	216,2±5	231±5
Anti-inflammatory (IC <sub>50</sub> values; µg/mL)					
RAW264,7	102±5	>400	82,04±5	95,4±5	

## Acknowledgments